

Original Paper

## Critical Review on Probiotics and its Effect on Cancer

Farzane Vafaeie \*

### ABSTRACT

Probiotics are defined as live microbial food ingredients that produce several beneficial effects to human health. The effects of probiotics may be classified in three modes of action. (i) Probiotics might be able to modulate the host's defenses. (ii) Probiotics can also have a direct effect on other microorganisms, commensal and/or pathogenic ones. (iii) Finally, probiotic effects may be based on actions affecting microbial products like toxins, host products e.g. bile salts and food ingredients. All three modes of probiotic action are in all likelihood involved in infection defense, prevention of cancer and in stabilizing or reconstituting the physiological balance between the intestinal microbiota and its host. Numerous bacteria in and on its external parts protect the human body from harmful threats. Several animal studies have shown that supplementation with specific strains of lactic acid bacteria (probiotics) could prevent the establishment, growth, and metastasis of transplantable and chemically induced tumors. An inverse relationship between the consumption of fermented dairy products, containing lactobacilli or bifidobacteria (are the main probiotic groups) *Pediococcus*, *Lactococcus*, *Bacillus*, *Shirota*, *Caseii*, *Lactis*, *Rhamnosus*, *Plantarum* and yeasts and the incidence of colon, gastric cancer and breast cancer has also been reported in epidemiological and population based case-control studies. The effectiveness in the treatment of cancers is based on the restoration of the sensitivity of transformed cells to apoptotic signals. The anticancer activity through induction of apoptosis of cancer cells seems to be promising approach for use of some probiotic strains as a support therapy or disease prevention. A wealth of data implicates that special receptors have essential roles in tumor development. A wealth of evidence emerging from laboratory studies indicates anticancer activity of probiotics.

**Keywords:** Probiotic, Anticancer Effect, Bifidobacterium, *Lactobacillus*, Receptor

### Author Information

\* Faculty of Science,  
Department of Biology,  
Ferdowsi University of  
Mashhad,  
Mashhad, Iran.

Submitted: 01.01.2016  
Accepted: 01.05.2016  
Published: 05.05.2016

One hundred years ago, we ate beneficial bacteria all the time but pasteurization, sterilization and irradiation of food have ended much of that. The concept of probiotics evolved at the turn of the twentieth century from a hypothesis first proposed by Nobel Prize-winning Russian scientist Elie Metchnikoff, who propounded that the long and healthy life of Bulgarian people resulted from their consumption of fermented milk products (1). Probiotic food can be defined as "food containing live microorganisms believed to actively enhance health by improving the balance of microflora in the

gut". A number of health benefits have been claimed for probiotic bacteria such as *Lactobacillus acidophilus*, *Bifidobacterium* spp., and *Lactobacillus casei*. Because of the potential health benefits, these organisms are increasingly incorporated into dairy foods (2).

### Molecular study

Increasingly, the microbiological scientific community is relying on molecular biology to define the complexity of the gut flora and to distinguish one organism from the next. Current techniques,

### References

1. Zhong W, Millsap K, Bialkowska-Hobrzanska H, Reid G. Differentiation of *Lactobacillus* species by molecular typing. *Applied and environmental microbiology*. 1998;64(7):2418-23.
2. Shahi N. *Probiotic bacteria: selective enumeration and survival in dairy foods*. *Journal of dairy science*. 2000;83(4):894-907.

Figure 1. The most commonly used species of lactic acid bacteria in probiotic preparations (3).

<i>Lactobacillus</i> sp.	<i>Bifidobacterium</i> sp.	<i>Enterococcus</i> sp.	<i>Streptococcus</i> sp.
<i>L. acidophilus</i>	<i>B. bifidum</i>	<i>Ent. faecalis</i>	<i>S. cremoris</i>
<i>L. casei</i>	<i>B. adolescentis</i>	<i>Ent. faecium</i>	<i>S. salivarius</i>
<i>L. delbrueckii</i> ssp. ( <i>bulgaricus</i> )	<i>B. animalis</i>		<i>S. diacetylactis</i>
<i>L. cellobiosus</i>	<i>B. infantis</i>		<i>S. intermedius</i>
<i>L. curvatus</i>	<i>B. thermophilum</i>		
<i>L. fermentum</i>	<i>B. longum</i>		
<i>L. lactis</i>			
<i>L. plantarum</i>			
<i>L. reuteri</i>			
<i>L. brevis</i>			

including genetic fingerprinting, gene sequencing, oligonucleotide probes and specific primer selection, discriminate closely related bacteria with varying degrees of success. Additional molecular methods being employed to determine the constituents of complex microbiota in this area of research are community analysis, denaturing gradient gel electrophoresis (DGGE)/temperature gradient gel electrophoresis (TGGE), fluorescent in situ hybridisation (FISH) and probe grids (4).

### Cancer

Cancer is a serious global public health problem. There are several epidemiological evidences that support a protective role of probiotics against cancer. A wealth of data implicates that ErbB receptors have essential roles in tumor development. Probiotic bacteria are known to exert an anticancer activity in animal studies(5).

Lactic acid bacteria and their probioactive cellular substances exert many beneficial effects in the gastrointestinal tract, and also release various enzymes into the intestinal lumen and exert potential synergistic (LAB) effects on digestion and alleviate symptoms of intestinal malabsorption (6). A carcinogen is any substance, radionuclide, or radiation that is an agent directly involved in causing cancer. This may be due to the ability to damage the genome or to the disruption of cellular metabolic processes.

### Colon Cancer

Colon cancer is the second to third most frequent type of cancer in Western industrialized countries. fiber diet, are of greater interest because they represent controllable risk factors. Within the complex gut microflora, which consists of >1 10<sup>11</sup>


### References

3. Parvez S, Malik K, Ah Kang S, Kim HY. Probiotics and their fermented food products are beneficial for health. *Journal of applied microbiology*. 2006;100(6):1171-85.
4. McCartney AL. Application of molecular biological methods for studying probiotics and the gut flora. *British Journal of Nutrition*. 2002;88(S1):s29-s37.
5. Ma EL, Choi YJ, Choi J, Pothoulakis C, Rhee SH, Im E. The anticancer effect of probiotic *Bacillus polyfermenticus* on human colon cancer cells is mediated through ErbB2 and ErbB3 inhibition. *International Journal of Cancer*. 2010;127(4):780-90.

living bacteria/g colon content, LAB belong to those bacteria with such beneficial effects (7) investigated for possible use as probiotics and for colon cancer biological products. Five of these strains inhibited

growth of eight food-borne pathogens including *Helicobacter pylori*, *Escherichia coli* and *Salmonella typhimurium* (8).

Figure 2. Mechanism of action of probiotics during colon cancer progression (6)



Several mechanisms have been proposed as to how LAB may inhibit colon cancer, including enhancing the host's immune response, altering the metabolic activity of the intestinal microbiota, binding and degrading carcinogens, producing antimutagenic compounds, and altering the physiochemical conditions in the colon (6) It appears that LAB can reduce the levels of colon enzymes that convert pro-carcinogens to carcinogens. Specifically, LAB can reduce levels of the enzymes  $\beta$ -glucuronides, nitro-reductase, and azo-reductase (Kampman et al. 1994).

Most animal and human studies do indicate that feeding certain LAB decreases fecal enzyme levels that may be involved in formation of carcinogens. Also, beneficial effects can be attributed to immune-potentiating effects by LAB strains. One specific effect was shown by heat-killed *Lactobacillus plantarum* L-137, which restored the inhibited IL-12 production in DBA/2 mice with tumors (9). The anti-tumor effects were found to be due to the activation of macrophage by LC9018 (10).

## References

6. Kumar M, Kumar A, Nagpal R, Mohania D, Behare P, Verma V, et al. Cancer-preventing attributes of probiotics: an update. *International journal of food sciences and nutrition*. 2010;61(5):473-96.
7. Wollowski I, Rechkemmer G, Pool-Zobel BL. Protective role of probiotics and prebiotics in colon cancer. *The American journal of clinical nutrition*. 2001;73(2):451s-5s.
8. Thirabunyanon M, Boonprasom P, Niamsup P. Probiotic potential of lactic acid bacteria isolated from fermented dairy milks on antiproliferation of colon cancer cells. *Biotechnology letters*. 2009;31(4):571-6.

### Breast Cancer

Results of cross-cultural and regional correlation studies have shown that breast cancer mortality correlates positively with the consumption of milk products (11). Milk fermented by *B. infantis*, *B. bifidum*, *B. animalis*, *L. acidophilus* and *L. paracasei* inhibited the growth of the MCF7 breast cancer cell line, and the antiproliferative effect was not related to the presence of bacteria (12). Isoflavones are a class of phytoestrogens plant-derived compounds with estrogenic activity. Soybeans and soy products are the richest sources of isoflavones in the human diet (13). Soy phytoestrogens were suggested to reduce the risk of a number of diseases including breast cancer. Given that these compounds are metabolized by bacteria, alteration of intestinal bacteria and enzymes may affect phytoestrogen metabolism (14). Observational studies suggest that dietary isoflavones reduce breast cancer risk, and this may be caused in part by effects on endogenous hormone concentrations. Because intestinal bacteria metabolize isoflavones, it was hypothesized that

consumption of probiotic bacteria would enhance the biologic effects of isoflavones, including effects on endogenous hormones (15).

### Bladder Cancers

Few human clinical trials conducted, one showed that *L. casei* consumption (three times per day for 1 year) increased the recurrence free period among subjects with bladder cancer (16).

Daily intake of a viable *L. casei* strain postponed recurrence of bladder tumors in a randomized, controlled, multicenter study in 48 Japanese patients. Patients were enrolled within 2 week after removal of one or more bladder tumors (17). One hypothesis for the prevention or delay of tumor development by lactobacilli is that they might bind to mutagenic compounds in the intestine (18).

Liver cancer: Liver cancer is the sixth most commonly diagnosed cancer in the world, and the third most common cause of death from cancer, according to Cancer Research UK (19).

### References

9. Murosaki S, Muroyama K, Yamamoto Y, Yoshikai Y. Antitumor effect of heat-killed *Lactobacillus plantarum* L-137 through restoration of impaired interleukin-12 production in tumor-bearing mice. *Cancer Immunology, Immunotherapy*. 2000;49(3):157-64.
10. Okawa T, Niibe H, Arai T, Sekiba K, Noda K, Takeuchi S, et al. Effect of LC9018 combined with radiation therapy on carcinoma of the uterine cervix. A phase III, multicenter, randomized, controlled study. *Cancer*. 1993;72(6):1949-54.
11. Armstrong B, Doll R. Environmental factors and cancer incidence and mortality in different countries, with special reference to dietary practices. *International Journal of Cancer*. 1975;15(4):617-31.
12. van't Veer P, Dekker JM, Lamers JW, Kok FJ, Schouten EG, Brants HA, et al. Consumption of fermented milk products and breast cancer: a case-control study in The Netherlands. *Cancer research*. 1989;49(14):4020-3.
13. Yildiz F. *Phytoestrogens in functional foods*: CRC Press; 2005.
14. Nettleton JA, Greany KA, Thomas W, Wangen KE, Adlercreutz H, Kurzer MS. Plasma phytoestrogens are not altered by probiotic consumption in postmenopausal women with and without a history of breast cancer. *The Journal of nutrition*. 2004;134(8):1998-2003.
15. Nettleton JA, Greany KA, Thomas W, Wangen KE, Adlercreutz H, Kurzer MS. Short-term soy and probiotic supplementation does not markedly affect concentrations of reproductive hormones in postmenopausal women with and without histories of breast cancer. *Journal of Alternative & Complementary Medicine: Research on Paradigm, Practice, and Policy*. 2005;11(6):1067-74.
16. De Roos NM, Katan MB. Effects of probiotic bacteria on diarrhea, lipid metabolism, and carcinogenesis: a review of papers published between 1988 and 1998. *The American journal of clinical nutrition*. 2000;71(2):405-11.
17. Orrhage K, Sillerström E, Gustafsson J-Å, Nord C, Rafter J. Binding of mutagenic heterocyclic amines by intestinal and lactic acid bacteria. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*. 1994;311(2):239-48.
18. El-Nezami HS, Polychronaki NN, Ma J, Zhu H, Ling W, Salminen EK, et al. Probiotic supplementation reduces a biomarker for increased risk of liver cancer in young men from Southern China. *The American journal of clinical nutrition*. 2006;83(5):1199-203.

Probiotics have also been shown to attenuate hepatotoxic effects of aflatoxin, a well known liver carcinogen, in rats (20) and to reduce biomarkers of liver cancer risk in a human intervention trial (20). Probiotic bacteria could block the intestinal absorption of aflatoxin B1 and thereby lead to reduced urinary excretion of aflatoxin B1-N7-guanine (AFB-N7-guanine), a marker for a biologically effective dose of aflatoxin exposure. Elevated urinary excretion of this aflatoxin-DNA adduct is associated with an increased risk of liver cancer (19).

---

### References

20. Gratz S, Täubel M, Juvonen R, Viluksela M, Turner P, Mykkänen H, et al. Lactobacillus rhamnosus strain GG modulates intestinal absorption, fecal excretion, and toxicity of aflatoxin B1 in rats. *Applied and environmental microbiology*. 2006;72(11):7398-400.

---

### Conclusion

The probiotic theory offers an intriguing approach to controlling negative metabolic or pathogenic activities of microbes to which we are exposed on a daily basis. Results suggest that consumption of a high amount of fermented milk products may have a protective effect on the risk of cancer (3).